

UNIT NO. 2

DIAGNOSIS OF DEMENTIA – HOW DO I KNOW THAT MY PATIENT HAS DEMENTIA?

Dr Chong Mei Sian

ABSTRACT

The prevalence of dementia is on a rising trend with the rapidly ageing population in Singapore. Hence, early diagnosis of dementia is important as early therapeutic interventions may palliate substantially, if not reverse, the significant emotional and economic costs of the illness.

In this paper, we present a 4-step approach to dementia evaluation, incorporating local data, where possible. The first step requires the exclusion of delirium as the cause of the forgetfulness or confusion. The second step involves establishing the diagnosis of dementia. The third step assesses the behavioural, functional and social problems associated with dementia. The final step, with the use of a focused history, physical examination, investigations and selected use of neuroimaging, attempts to establish the aetiological diagnosis of the dementia.

INTRODUCTION

With the rising trend in the prevalence of dementia, especially with Singapore's rapidly greying population, this is an area of intense research in the area of therapeutics as well as the diagnosis of dementia in the earlier stages, even in the preclinical dementia state as with early diagnosis, the affected patients are then more amenable to treatment advances.

MILD COGNITIVE IMPAIRMENT

Cognitive changes in the elderly occur over a continuum, ranging from normal ageing at one end of the spectrum to dementia at the other end. There has been intense interest in the intermediate stage between normal ageing and dementia. Of the various classification systems, the Mayo Clinic's mild cognitive impairment (MCI) has received the most attention. Its pathological validity is supported by conversion rates to dementia of approximately 12% annually and 80% at six years of follow-up. Originally, MCI diagnosis required the presence of memory complaint (preferably corroborated by an informant), objective memory impairment for age, essentially preserved general cognitive function, normal functional activities and no dementia. The heterogeneity within MCI has led to the proposal of a new classification system, based predominantly on neuropsychological profiles and includes amnesic or single memory MCI, multiple-domain MCI and single non-memory MCI. However, there are still many controversies surrounding

this clinical entity and initial drug trials have not shown clinical benefit. As such, the discussion below will focus mainly on established dementia.

ASSESSMENT

The evaluation of dementia should be targeted at individuals in whom there is some suspicion of cognitive impairment. This includes subjects with memory or other cognitive complaints, this could either be self-reported or noticed by family members or caregivers; subjects in whom the physician has suspicion of cognitive impairment during the consultation despite the absence of memory or cognitive complaints; subjects who are at increased risk for dementia, such as those with strong family history of dementia and elderly subjects who need to make an important decision (such as making a will, sale of flat, handling complicated financial matters) and in whom mental competency is in question.

The evaluation of cognitive impairment should have been done via a multifaceted approach, focusing not only on the cognitive complaints, but also on the functional and social consequences of these cognitive changes. This would help the clinician diagnose dementia early, assess for the complications of dementia and establish the aetiology of the dementia and manage accordingly.

With a patient presenting with forgetfulness or confusion, we can use a 4-step assessment to evaluate the cognitive complaint:

- (i) Is the forgetfulness or confusion acute or chronic?
- (ii) If the forgetfulness or confusion is chronic, is it dementia?
- (iii) If it is dementia, what are the complications?
- (iv) If it is dementia, what is the aetiology?

(i) Is the Forgetfulness or Confusion Acute or Chronic?

If the cognitive complaints is of an acute nature, with a rapid onset and short duration (lasting from few hours to days), it would be important to exclude delirium.

Delirium is defined by the Diagnostic and Statistical Manual of Mental Disorders – fourth edition (DSM-IV), however, this may be difficult to apply in clinical practice. The Confusion Assessment Method (CAM) is a brief and structured assessment commonly used in clinical setting to diagnose delirium. It requires the presence of 3 of the following 4 features: presence of acute change in mental status and fluctuating course with inattention, coupled with either the presence of disorganised thinking or altered level of consciousness. CAM has been shown to have 94-100% sensitivity and 90-100% specificity in the identification of delirium with good inter-observer reliability (kappa test 0.81-1.0). If the cognitive complaints are assessed to be secondary to delirium, the underlying precipitating factors

(such as sepsis, stroke disease or drug causes) should be looked out for and the patient would require hospitalisation to manage the delirium and the underlying medical illness.

One must also be mindful that acute confusional state can sometimes be superimposed on chronic confusion. If the forgetfulness or confusion is of a subacute nature, developing over a period of week to few months, conditions such as stroke disease, space-occupying lesion, Creutzfeld-Jakob disease and hydrocephalus have to be excluded.

(ii) Is It Dementia?

If the cognitive complaints are of a chronic nature, it is first important to exclude depression and late-onset psychiatry disorders. The diagnosis of dementia is then assessed via a clinical approach, either subjectively (looking for features of cognitive decline in the subject) or objectively (testing the subject's cognitive abilities using validated performance-based assessments).

Subjective Approach

The DSM-IV criteria for dementia are often used as the gold standard for clinical diagnosis of dementia. It requires the presence of memory impairment, together with deficits in one other cognitive domain (aphasia, apraxia, agnosia and executive dysfunctioning). Examples of practical questions to be asked of the patient's informants with regards to these cognitive domains are shown in Table 1.

The Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) is a 26-item test that enquires about the subject's memory, cognition and language ability for the last 10 years. The strength of this instrument is its assessment of cognitive changes over a period of time, instead of a single point in time and also it is independent of the subject's premorbid ability or past educational attainments. This has also been validated locally among elderly Chinese subjects with an optimal cut-off score of 3.3/3.4 with 94.3% sensitivity and 94% specificity.

Objective Approach

This is an observer-based approach using either performance-based instruments, such as mental status test (brief screening instruments), or a more detailed neuropsychological tests, which is usually administered by clinical psychologists.

There are several mental status tests which have been validated locally. These include the Elderly Cognitive Assessment Questionnaire (ECAQ), Abbreviated Mental Test (AMT) and the Chinese Mini Mental Status Examination (CMMSE).

The ECAQ is a 10-item cognitive test which assesses memory and information-orientation. Using a cut-off score of 5/6, it has 85.3% sensitivity with 91.5% specificity for identifying cognitive impairment.

The 10-item AMT (Table 2) and 28-item CMMSE have also been validated locally. For mild cognitive impairment, AMT's cut-off score is 7/8 (81% sensitivity with 89% specificity) and CMMSE cut-off score of 20/21 (sensitivity 83%, specificity

Table 1. DSM-IV Clinical Criteria for Diagnosis of Dementia

Cognitive domain	Questions
Amnesia	Any forgetfulness? Did it start gradually or suddenly? Is it progressively worse? And if so, is it smoothly declining or showing a step-wise/fluctuating decline? Is it over short-term or long-term matters?
AND declines in one of the following domains:	
Aphasia	Any word-finding difficulty or other difficulties with communication?
Apraxia	Any problems with buttoning or dressing? Any difficulties with using utensils during mealtimes?
Agnosia	Any problems recognising familiar faces or familiar items?
Executive dysfunctioning	Any problems handling money (loose change)? Any change in general problem-solving abilities? Is one's work getting to be more disorganised?
OF sufficient severity to cause significant impairment in social or occupational functioning	As a result of the above, is he becoming less independent in the - community? - home-care? - self-care level?

Table 2. Abbreviated Mental Test (AMT)

Items	Score
What is the year?	1
What is the time? (within 1 hour)	1
What is your age?	1
What is your date of birth?	1
What is your home address?	1
Where are we now?	1
Who is our country's Prime Minister?	1
What is his/her job? (show picture)	1
Memory phrase "37 Bukit Timah Road"	
Count backwards from 20 to 1	1
Recall memory phrase	1
Total score	

94%). The CMMSE is more useful in those with higher educational attainment as the AMT has a ceiling effect on these individuals.

Neuropsychological testing is useful in detecting subtle cognitive difficulties which are not picked up by the brief screening instruments. They should be performed on subjects who have memory complaints but do not yet satisfy criteria for dementia; depressed subjects who present with memory complaints to help in determining whether the memory complaints are due solely to the depression or whether they have concomitant dementia; and subjects in whom decision-making capacity is being assessed. Psychometric testing can be a useful adjunct in the latter scenario. Neuropsychometric batteries have been validated locally in the elderly Chinese and the Vascular Dementia Battery test has also been validated in the Singapore population.

Neuropsychological tests are also useful in individuals in whom the diagnosis of dementia is inconclusive and serial monitoring for performance decline over time is useful in establishing the diagnosis.

(iii) What Are the Dementia Complications?

The complications of dementia can be broadly divided into behavioural and psychological symptoms (discussed in Unit 5), functional problems and social problems (discussed in Units 3 and 6). These should be evaluated in all patients with dementia as these issues are the major causes of stress on the caregiver and assessment would enable the clinician to target subsequent management effectively.

Functional difficulties can be assessed at 3 levels: community functioning, home functioning and self-care. They are generally affected with the progression of dementia in a descending order and also allow these functional deficits to serve as markers of dementia severity. It is important when asking for functional deficits to ask for a change in the level of function, i.e. whether the patient is functioning at the same level as before and whether the patient is as independent as before. It is also important to make sure that these difficulties result from cognitive difficulties and not physical disabilities.

The severity of dementia can be staged using the Diagnostic and Statistical Manual of Mental Disorders-3rd revised-edition (DSM-III-R) criteria where mild dementia is defined as impairment for work and social activities with the capacity for independent living remaining largely intact. Moderate dementia takes place when independent living is hazardous and would require some degree of supervision. Severe dementia is characterized by impaired activities of daily living such that continual supervision is required. Other formal functional assessment scales include Clinical Dementia Rating Scale (CDR), Functional Assessment Staging (FAST), Barthel Index and Blessed Dementia Scale (BDS).

(iv) What Is the Dementia Aetiology?

Having determined the cognitive impairment to be chronic and having met clinical criteria for dementia, as well as assessing for the complications of dementia, the final step of the clinical evaluation involves determining the dementia aetiology.

The types of dementia can be broadly divided into 2 categories – irreversible and reversible causes (Table 3). The aim of determining dementia aetiology is to rule out potentially reversible causes of dementia and selecting appropriate treatment strategies for the irreversible dementias. This is done via clinical history and physical examination, followed by laboratory investigations and neuroimaging. There are guidelines and

practice parameters developed for evaluating of dementia etiology and also more specific criteria for diagnosis of the more common Alzheimer's disease (AD) and vascular dementia (VD).

In the history, it is important to ask for the nature of the cognitive decline (sudden or gradual), progression – either gradually progressive (more suggestive of AD) or stepwise/fluctuating course (suggestive of VD). A history of significant alcohol ingestion and medication use (such as antipsychotics, antidepressants, anticholinergic agents and sedative-hypnotic agents) and history of medical, neurological and psychiatric illness is important.

A targeted physical examination should be performed, looking for focal neurological deficits (such as visual field defects, hemiparesis, hemisensory loss, asymmetric deep tendon reflexes or unilateral extensor plantar responses). It is also important to examine for extrapyramidal signs such as rigidity and bradykinesia, movement disorders and gait abnormalities as these may point to certain aetiological diagnosis.

Dementias which are related to metabolic abnormalities are thought to be reversible. The most commonly recommended haematological tests are: full blood count, urea and electrolytes, serum calcium, serum glucose, thyroid function tests and vitamin B12 levels. We do not advise routine testing for neurosyphilis given the problems in interpreting the results of testing. Serum Venereal Disease Research Laboratory (VDRL) testing detects only 75% of tertiary syphilis and CSF VDRL may be negative in 30-70% of cases and neurosyphilis. Thus we recommend testing only when patients exhibit clinical features of neurosyphilis.

Other biomarkers which can help in establishing dementia diagnosis include apolipoprotein-E e4 allele, CSF-tau and β -amyloid for AD, CSF 14-3-3, neuron-specific enolase and electroencephalogram for Creutzfeldt-Jakob disease. However, these are not performed routinely.

Neuroimaging is useful in the differential diagnosis of dementia and also necessary in the diagnostic criteria in AD and VD. It is also useful in detection of very early dementia as the functional and structural brain changes takes place before clinical manifestation of cognitive deficits. It consists of either structural imaging techniques (computed tomography (CT) scan of head and magnetic resonance imaging (MRI)) or functional neuroimaging techniques (Positron emission tomography and single-photon emission tomography). Whether all patients with dementia require a structural imaging is an important clinical question, for which there is no consensus. The value of neuroimaging is the identification of cerebral infarcts and clinically important surgical brain lesions (SBLs) such as subdural haematomas, cerebral tumors and normal pressure hydrocephalus. The Canadian Consensus Conference on the Assessment of Dementia (CCCAD) has outlined the criteria for undertaking a CT scan, only if certain conditions are met (Table 4). We also believe that the functional stage of the dementia is also relevant and important, over and above the duration of cognitive symptoms. In a patient with advanced dementia of long duration (>2 years), we believe that a brain scan is not warranted to detect potentially reversible SBLs.

Table 3. Types of Dementia

Irreversible causes

- o Degenerative causes – Alzheimer's disease (AD), frontotemporal dementia, diffuse Lewy body dementia
- o Cerebrovascular disease – vascular dementia (VD)
- o Prion-associated disorders (Creutzfeldt-Jakob disease)
- o Neurogenetic disorders

Potentially reversible causes

- o Infectious disorders – meningitis, encephalitis
- o Toxic or metabolic causes – hypothyroidism, vitamin B12 deficiency, alcohol-related syndromes
- o Neoplastic causes
- o Hydrocephalus – obstructive or normal pressure hydrocephalus

Table 4. Canadian Consensus Conference Criteria for Performing Cranial CT in Patients with Dementia

- o Patients are less than 60 years old
- o Rapid (e.g. over 1-2 months), unexplained decline in cognition or function
- o Dementia of relatively short duration (< 2 yrs)
- o Recent, significant head trauma
- o Unexplained neurologic symptoms (e.g. new onset of severe headache or seizures)
- o History of cancer, especially of a type or at a site associated with metastasis to the brain
- o Use of anticoagulants or history of bleeding disorder
- o History of urinary incontinence and gait disturbance early in the course of dementia (suggestive of normal pressure hydrocephalus)
- o Presence of any new localizing signs on physical examination (hemiparesis, babinski's sign)
- o Unusual or atypical cognitive symptoms or presentation (e.g. progressive aphasia)
- o Gait disturbance

CT is recommended if one or more of these criteria are present.

However, if the patient's dementia is still mild and moderate (even after 2 years), it would be advisable to request for an initial CT brain. When caregivers with advanced dementia patients present to the clinic, most of the time, they are seeking help for the management of the behavioural, functional and social problems and not in the precise aetiological diagnosis. However, it would be imperative to discuss with the caregivers and seek their agreement if the clinician is not inclined to perform neuroimaging based on the above discussion and the CCCAD criteria.

Now, having completed the clinical evaluation and ruled out potentially reversible causes of dementia, many clinical criteria for the two most common types of dementia (AD and VD) have been developed over the years. In AD, the common clinical criteria used include DSM-IV criteria for DAT, National Institute of Neurologic, Communicative Disorders and Stroke-AD and Related Disorders Association (NINCDS-ADRDA). In VD, the common clinical criteria used include DSM-III-R, National Institute of Neurologic Disorder and Stroke and the Association Internationale pour la Recherche et l'Enseignement

en Neurosciences (NINDS-AIREN), State of California AD Diagnostic and Treatment Centers criteria (ADDTCC), Hachinski Ischemic Score (HIS), modified Rosen scale (the latter two being clinical scales). These can be used in the specialized dementia clinics for the definition of AD and VD.

CONCLUSIONS

In summary, a 4-step clinical approach is useful in evaluating the individual who presents with forgetfulness or confusion. This is summarized succinctly in the summary table below which can aid the family physician when a patient presents to clinic with cognitive complaints.

Table 5. Summary of Approach to Patient with Memory Complaint

- o Is the memory complaint acute or chronic?
Rule out delirium
- o If it is chronic, is it dementia ?
- o If it is dementia, what are the complications?
Behavioural, functional, social aspects of dementia
- o What is the aetiology?
Clinical evaluation (history, clinical examination, laboratory tests, + neuroimaging)
To rule out reversible causes.
If irreversible cause, clinical criteria in the differential diagnosis of dementia aetiology.

RECOMMENDED READING

1. MOH Clinical Practice Guidelines. September 2001. no 5/01.
2. MS Chong, S Sahadevan. An Evidence-based Clinical Approach to the diagnosis of dementia. Annals Academy of Medicine Singapore 2003; 32:740-8.
3. Knopman DS, DeKosky ST, Cummings JL, Chui H, Corey-Bloom J, Relkin N, Small GW, Miller B, Stevens JC. Practice parameter: Diagnosis of dementia (an evidence-based review) Report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology 2001; 56:1143-53.
4. Peterson RC. Conceptual overview. IN: Petersen RC, ed. Mild cognitive impairment: Aging to Alzheimer's disease. New York, NY: Oxford University Press, Inc; 2003:1-14.

LEARNING POINTS

- o **Early diagnosis of dementia is important as early therapeutic interventions may palliate substantially, if not reverse, the significant emotional and economic costs of the illness.**
- o **A four-step method can be used.**
- o **The first step requires the exclusion of delirium as the cause of the forgetfulness or confusion.**
- o **The second step involves establishing the diagnosis of dementia.**
- o **The third step assesses the behavioural, functional and social problems associated with dementia.**
- o **The final step attempts to establish the aetiological diagnosis of the dementia.**